

Preventing Intimate Partner Violence

What is intimate partner violence?

Intimate partner violence (IPV) is abuse or aggression that occurs in a romantic relationship. “Intimate partner” refers to both current and former spouses and dating partners. IPV can vary in how often it happens and how severe it is. It can range from one episode of violence that could have lasting impact to chronic and severe episodes over multiple years.

IPV can include any of the following types of behavior:¹

- **Physical violence** is when a person hurts or tries to hurt a partner by hitting, kicking, or using another type of physical force.
- **Sexual violence** is forcing or attempting to force a partner to take part in a sex act, sexual touching, or a non-physical sexual event (e.g., sexting) when the partner does not or cannot consent.
- **Stalking** is a pattern of repeated, unwanted attention and contact by a partner that causes fear or concern for one’s own safety or the safety of someone close to the victim.
- **Psychological aggression** is the use of verbal and non-verbal communication with the intent to harm a partner mentally or emotionally and/or to exert control over a partner.



IPV is connected to other forms of violence and is related to serious health issues and economic consequences. However, IPV and other forms of violence can be prevented.

How big is the problem?

IPV is common. It affects millions of people in the United States each year. Data from CDC’s National Intimate Partner and Sexual Violence Survey (NISVS) indicate:²

- About 41% of women and 26% of men experienced contact sexual violence, physical violence, and/or stalking by an intimate partner and reported an intimate partner violence-related impact during their lifetime. Injury, posttraumatic stress disorder (PTSD) symptoms, concern for safety, fear, needing help from law enforcement, and missing at least one day of work are common impacts reported.
- Over 61 million women and 53 million men have experienced psychological aggression by an intimate partner in their lifetime.

IPV starts early and continues throughout people’s lives. When IPV occurs in adolescence, it is called teen dating violence (TDV). TDV affects millions of U.S. teens each year. About 16 million women and 11 million men who reported experiencing contact sexual violence, physical violence, or stalking by an intimate partner in their lifetime said that they first experienced these forms of violence before the age of 18.² While violence impacts all people in the United States, some individuals and communities experience inequities in risk for violence due to the social and structural conditions in which they live, work and play. Youth from groups that have been marginalized, such as sexual and gender minority youth, are at greater risk of experiencing sexual and physical dating violence.^{3,4}



Centers for Disease Control and Prevention
National Center for Injury Prevention and Control

What are the consequences?

IPV is a significant public health issue that has many individual and societal costs. About 75% of female IPV survivors and 48% of male IPV survivors experience some form of injury related to IPV.² IPV can also result in death. Data from U.S. crime reports suggest that about 1 in 5 homicide victims are killed by an intimate partner. The reports also found that over half of female homicide victims in the United States are killed by a current or former male intimate partner.⁶

Many other negative health outcomes are associated with IPV. These include a range of conditions affecting the heart, muscles and bones, and digestive, reproductive, and nervous systems, many of which are chronic.⁵ Survivors can experience mental health problems such as depression and PTSD symptoms. They are at higher risk for engaging in behaviors such as smoking, binge drinking, and sexual risk activity.⁷ People from groups that have been marginalized, such as people from racial and ethnic minority groups, are at higher risk for worse consequences.⁸

Although the personal consequences of IPV are devastating, there are also many costs to society. The lifetime economic cost associated with medical services for IPV-related injuries, lost productivity from paid work, criminal justice, and other costs is \$3.6 trillion. The cost of IPV over a victim's lifetime was \$103,767 for women and \$23,414 for men.⁹

How can we prevent intimate partner violence?

Promoting healthy, respectful, and nonviolent relationships and communities can help reduce the occurrence of IPV. It also can prevent the harmful and long-lasting effects of IPV on individuals, families, and communities. CDC developed a resource, *Preventing Intimate Partner Violence Across the Lifespan: A Technical Package of Programs, Policies, and Practices*, to help communities use the best available evidence to prevent intimate partner violence. This resource can be used as a tool in efforts to impact individual behaviors as well as family, community, and society factors that influence risk and protective factors for intimate partner violence.⁷

Preventing Intimate Partner Violence Across the Lifespan:

A Technical Package of Programs, Policies, and Practices

A **technical package** is a collection of strategies based on the best available evidence to prevent or reduce public health problems. The **strategy** lays out the direction and actions to prevent intimate partner violence. The **approach** includes the specific ways to advance the strategy through programs, policies and practices. The **evidence** for each of the approaches in preventing intimate partner violence and associated risk factors is also included.

References

1. Breiding, M. J., Basile, K. C., Smith, S. G., Black, M. C., & Mahendra, R. R. (2015). Intimate partner violence surveillance: uniform definitions and recommended data elements, Version 2.0. Atlanta (GA): National Center for Injury Prevention and Control, Centers for Disease Control and Prevention.
2. Leemis R.W., Friar N., Khatiwada S., Chen M.S., Kresnow M., Smith S.G., Caslin, S., & Basile, K.C. (2022). The National Intimate Partner and Sexual Violence Survey: 2016/2017 Report on Intimate Partner Violence. Atlanta, GA: National Center for Injury Prevention and Control, Centers for Disease Control and Prevention.
3. Johns MM, Lowry R, Andrzejewski J, Barrios LC, Demissie Z, McManus T, Rasberry CN, Robin L, Underwood JM. Transgender identity and experiences of violence victimization, substance use, suicide risk, and sexual risk behaviors among high school students—19 states and large urban school districts, 2017. *Morbidity and Mortality Weekly Report*. 2019 Jan 25;68(3):67.
4. Johns MM, Lowry R, Haderkhanaj LT, Rasberry CN, Robin L, Scales L, Stone D, Suarez NA. Trends in violence victimization and suicide risk by sexual identity among high school students—Youth Risk Behavior Survey, United States, 2015–2019. *MMWR supplements*. 2020 Aug 21;69(1):19.
5. Smith, S.G., Chen, J., Basile, K.C., Gilbert, L.K., Merrick, M.T., Patel, N., Walling, M., & Jain, A. (2017). The National Intimate Partner and Sexual Violence Survey (NISVS): 2010–2012 State Report. Atlanta, GA: National Center for Injury Prevention and Control, Centers for Disease Control and Prevention.
6. Jack SP, Petrosky E, Lyons BH, et al. Surveillance for Violent Deaths — National Violent Death Reporting System, 27 States, 2015. *MMWR Surveill Summ* 2018;67(No. SS-11):1–32. DOI: <http://dx.doi.org/10.15585/mmwr.ss6711a1external>
7. Niolon, P. H., Kearns, M., Dills, J., Rambo, K., Irving, S., Armstead, T., & Gilbert, L. (2017). Preventing Intimate Partner Violence Across the Lifespan: A Technical Package of Programs, Policies and Practices. Atlanta, GA: National Center for Injury Prevention and Control, Centers for Disease Control and Prevention.
8. Stockman JK, Hayashi H, Campbell JC. Intimate Partner Violence and its Health Impact on Ethnic Minority Women [corrected] [published correction appears in *J Womens Health (Larchmt)*. 2015 Mar;24(3):256]. *J Womens Health (Larchmt)*. 2015;24(1):62-79. doi:10.1089/jwh.2014.4879
9. Peterson, C., Kearns, M. C., McIntosh, W. L., Estefan, L. F., Nicolaidis, C., McCollister, K. E., & Florence, C. (2018). Lifetime Economic Burden of Intimate Partner Violence Among U.S. Adults. *American Journal of Preventive Medicine*, 55(4), 433–444.



Teach safe and healthy relationship skills

- Social-emotional learning programs for youth
- Healthy relationship programs for couples



Engage influential adults and peers

- Men and boys as allies in prevention
- Bystander empowerment and education
- Family-based programs



Disrupt the developmental pathways toward partner violence

- Early childhood home visitation
- Preschool enrichment with family engagement
- Parenting skill and family relationship programs
- Treatment for at-risk children, youth, and families



Create protective environments

- Improve school climate and safety
- Improve organizational policies and workplace climate
- Modify the physical and social environments of neighborhoods



Strengthen economic supports for families

- Strengthen household financial security
- Strengthen work-family supports



Support survivors to increase safety and lessen harms

- Victim-centered services
- Housing programs
- First responder and civil legal protections
- Patient-centered approaches
- Treatment and support for survivors of IPV, including teen dating violence